

The Centers for Disease Control and Prevention's Laboratory Outreach and Communication System (LOCS) is forwarding this official Health Advisory "Influenza-Associated Pediatric Mortality and the Increase of *Staphylococcus aureus* co-infection." As stated in the advisory, clinicians, clinical agencies and medical examiners are being asked to notify their local or state health department as soon as possible when deaths among children due to laboratory-confirmed influenza are identified. Clinical laboratories should be aware of possible requests from local or state health departments for *S. aureus* isolates from pediatric influenza patients in whom co-infection was noted. Because it is possible that cultures could be discarded before a request from local or state health departments is communicated to the laboratory, it is important for clinical laboratories to maintain *S. aureus* isolates from sterile sites/sputum/endotracheal cultures for future requests. Please disseminate this information through your most appropriate channels.

This is an official CDC Health Advisory distributed via the Health Alert Network (HAN) on May 9, 2007, 11:15 EDT (11:15 AM EDT) .

## **CDC Health Advisory**

### **Influenza-Associated Pediatric Mortality and the Increase of *Staphylococcus aureus* co-infection**

CDC is requesting that states report all cases of influenza-related pediatric mortality from the 2006-2007 influenza season.

Since 2004, the Influenza-Associated Pediatric Mortality Surveillance System, part of the Nationally Notifiable Disease Surveillance System, has collected information on deaths among children due to laboratory-confirmed influenza, including the presence of other medical conditions and bacterial infections at the time of death. From October 1, 2006 through May 7, 2007, 55 deaths from influenza in children have been reported to CDC from 23 state health departments and two city health departments. Data on bacterial co-infections were reported for 51 cases; 20 (39%) had a bacterial co-infection, and 16/20 were infected with *Staphylococcus aureus*. While the number of pediatric influenza associated deaths is similar to that reported during the two previous years, there has been an increase in the number of deaths in which both influenza and pneumonia or bacteremia due to *S. aureus* were identified. Only one influenza and *S. aureus* co-infection was identified in 2004-2005, and 3 were identified in 2005-2006. Of the 16 children reported with *S. aureus* so far in 2006-2007, 11 children had methicillin-resistant (MRSA) isolated from a sterile site (9) or sputum (2), and 5 had methicillin-susceptible *S. aureus* isolated from a sterile site (3) or sputum (2). The median age of children with *S. aureus* co-infection was older than children without *S. aureus* co-infection (11 years versus 4 years,  $p < .01$ ). Children with influenza and *S. aureus* co-infections were reported to be in good health before illness onset but progressed rapidly to severe illness. Influenza strains isolated from these children have not been different from common strains circulating in the community and the MRSA strains have been typical of those associated with MRSA skin infection outbreaks in the United States.

Healthcare providers should be alerted to the possibility of bacterial co-infection among children with influenza, and request bacterial cultures when bacterial co-infection is

suspected. Clinicians, clinical agencies and medical examiners are asked to contact their local or state health department as soon as possible when deaths among children due to laboratory-confirmed influenza are identified. CDC requests that all cases of pediatric influenza-associated deaths be reported promptly by state health departments to CDC through <http://sdn.cdc.gov> and that information about bacterial pathogens isolated from sterile sites and/or from sputum or endotracheal aspirates be completed on the Influenza-Associated Pediatric Mortality Surveillance System case report form. If the influenza death was complicated by *S. aureus* infection, please contact the clinical agency to determine if the *S. aureus* isolate is available. CDC is interested in receiving *S. aureus* isolates to better characterize those from fatal cases of influenza in children.

If you have any questions about this Health Advisory, please call the Influenza Division, Epidemiology and Prevention Branch at 404-639-3727.

Categories of Health Alert messages:

**Health Alert** conveys the highest level of importance; warrants immediate action or attention.

**Health Advisory** provides important information for a specific incident or situation; may not require immediate action.

**Health Update** provides updated information regarding an incident or situation; unlikely to require immediate action.

##This Message was distributed to State Epidemiologists##

=====



*Links to non-Federal organizations are provided solely as a service to our users. Links do not constitute an endorsement of any organization by CDC or the Federal Government, and none should be inferred. The CDC is not responsible for the content of the individual organization Web pages found at this link.*

Categories of LOCS messages:

**LOCS Alert** - conveys the highest level of importance; warrants immediate action or attention.

**LOCS Advisory** - provides important information for a specific incident or situation; may not require immediate action.

**LOCS Update** - provides updated information regarding an incident or situation; unlikely to require immediate action.